

#### WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# NTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

nternational Patent Classification  $^7$ :

C07F 9/38, A61K 31/663, 47/26, 9/48, 9/12

(11) International Publication Number: A1

WO 00/34293

(43) International Publication Date:

15 June 2000 (15.06.00)

(21) International Application Number:

PCT/US99/29284

(22) International Filing Date:

10 December 1999 (10.12.99)

(30) Priority Data:

09/209,153 ¥09/414,401 09/456,460

10 December 1998 (10.12.98) 7 October 1999 (07.10.99) 8 December 1999 (08.12.99)

US US US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications

Filed on

09/209,153 (CIP) 10 December 1998 (10.12.98) 09/414,401 (CIP)

US Filed on US Filed on

7 October 1999 (07.10.99) 09/456,460 (CIP) 8 December 1999 (08.12.99)

(71) Applicant (for all designated States except US): AESGEN, INC. [US/US]; P.O. Box 7275, Priceton, NJ 08543-7275 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): SHINAL, Edward, C. [US/US]; 10 Sherwood Court, Holmdel, NJ 07733 (US).

(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).

(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

**Published** 

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: METHOD FOR PREPARATION OF DISODIUM PAMIDRONATE


(57) Abstract

The invention provides a method of preparing anhydrous disodium pamidronate or an aqueous disodium pamidronate solution for pharmaceutical use.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL AM AT AU AZ BA BB BE BF BG BJ BR CCA CCF CCG	Albania Armenia Austria Austria Azerbaijan Bosnia and Herzegovina Barbados Belgium Burkina Faso Bulgaria Benin Brazil Belarus Canada Central African Republic Congo	FI FR GA GB GE GN GR HU IE IL IS IT JP KE	Spain Finland France Gabon United Kingdom Georgia Ghana Guinea Greece Hungary Ireland Israel Iceland Italy Japan Kenya	LT LU LV MC MD MG MK ML MN MN MN MR MV MX NE NL	Lithuania Luxembourg Larvia Monaco Republic of Moldova Madagascar The former Yugoslav Republic of Macedonia Mali Mongolis Mauriania Malawi Mexico Niger Netherlands	SK SN SZ TD TG TJ TM TR TT UA UG US UZ VN YU ZW	Slovakia Senegal Swaziland Chad Togo Tajikistan Turkmenistan Turkey Traidad and Tobago Ulcraine Uganda United States of America Uzbekistan Viet Nam Yugoslavia Zimbabwe
		GR					
	•		<b>4</b> •	MN	Mongolis	UA	Ukraine
BR	Brazil			*			•
		IT		MX	Mexico		
		_	•		_	YU	Yugoslavia
СН	Switzerland	KG KP	Kyrgyzstan Democratic People's	NO NZ	Norway New Zealand	ZW	Zmbabwe
CI	Côte d'Ivoire Cameroon	KF	Republic of Korea	PL	Poland		
CN	China Cuba	KR K2	Republic of Korea Kazakstan	PT RO	Portugal Romania		
CZ	Czech Republic	LC	Saint Lucia	RU SD	Russian Federation Sudan		
DE DK	Germany Denmark	u LK	Liechtenstein Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



20

#### METHOD FOR PREPARATION OF DISODIUM PAMIDRONATE

5 This application is a continuation-in-part of U.S. Patent Application Serial No. 09/414401, filed October 7, 1999 which is a continuation-in-part of U.S. Patent Application Serial No. 09/209153, filed December 10, 1998.

#### Background of the Invention

Disodium 3-amino-1-hydroxypropane-1,1-diphosphonate or "disodium 10 pamidronate" has the formula:

$$\begin{array}{c} \text{HO-P(O)--ONa} \\ \text{H}_2\text{N}-\text{CH}_2\text{CH}_2-\text{C}-\text{OH} \\ \text{HO-P(O)--ONa} \end{array}$$

It is commercially available as the lyophilized pentahydrate, under the name AREDIA® from Novartis Pharmaceuticals Corp., and is used to inhibit bone resorption, i.e., to treat moderate or severe hypercalcemia associated with malignancy, with or without bone metastases. The preparation of the crystalline pentahydrate from pamidronic acid is disclosed by Stahl et al. in U.S. Pat. Nos. 4,711,880 and 4,639,338. This material is prepared by partially neutralizing a heated slurry of pamidronic acid with aqueous sodium hydroxide (NaOH) to pH 7-7.5 and then initiating crystallization at ≥ 50°C. The mixture is slowly cooled to 0-5°C and the disodium pamidronate is collected by filtration. This product is described as having "excellent crystallinity" and as being "completely stable to storage under approximately normal ambient conditions." It comprises about 24.1-25% water.

The crystalline product ("Modification E") is contrasted with an

25 amorphous product disclosed to be prepared by the general process of Jary et al.

(U.S. Pat. No. 4,304,734). In Comparison Example 1 of the '338 patent,

pamidronic acid is neutralized to pH 7.4 with aqueous NaOH and the reaction

mixture is concentrated to dryness under reduced pressure at 60-70°C and then

dried at 20 mbar to constant weight. However, the amorphous product that is

30 obtained is described as deliquescent in air, contains 12.9% of retained water

and, as taught in the '338 patent, will convert to the pentahydrate. Stahl et al.



10

15

20

also teach the interconversion of other crystalline forms of disodium pamidronate depending upon humidity and amount of water present, which demonstrates the difficulty in utilizing preformed disodium salts of pamidronic acid for further processing into sterile pharmaceutical dosage forms.

Thus, a need exists for a simple method to prepare disodium pamidronate and dosage forms thereof that are stable to storage and suitable for reconstitution and/or intravenous infusion/injection.


#### Summary of the Invention

The present invention provides a pharmaceutical unit dosage form comprising an aqueous solution of disodium pamidronate enclosed in a suitable container, such as an ampule or vial, under an inert atmosphere. Preferably the container is designed for "single use". Preferably, the container is formed of a material that does not contain calcium in a form that can be sequestered by the issolved disodium pamidronate.

The present invention also provides a method for the preparation of amorphous, essentially anhydrous ( $\leq 1-2$  wt-% water) disodium pamidronate (the "product"). This solid product can be stored under nitrogen at otherwise ambient conditions, and readily reconstituted with sterile water or physiological salt solutions for injection or infusion into a patient in need of treatment therewith.

The present method also comprises a method to make an aqueous solution of disodium pamidronate by addition of aqueous sodium nyuroxide to a stirred slurry of pamidronic acid in an about 2:1 molar ratio of NaOH to acid, optionally comprising excess mannitol, to yield a clear solution of pH  $6.5 \pm 0.1$ . Aqueous phosphoric acid is added as needed if the pH following NaOH addition is too high. A small amount of aqueous NaOH can be added if the pH is too low. Preferably, 1 N aqueous NaOH is used.

The solution is frozen and lyophilized under reduced pressure to yield amorphous, essentially anhydrous disodium pamidronate. The product is stable when stored under dry nitrogen at otherwise ambient conditions. Preferably, the about pH 6.5 solution is filtered and aliquots are introduced into suitable container(s), then frozen (-25 to -40°C), lyophilized (10-25 mbar, 20-40°C) in sinu, and the containers sealed under positive nitrogen pressure to yield a



25

plurality of unit dosage forms of solid disodium pamidronate, optionally in admixture with mannitol.

For example, vials containing 30 mg, 60 mg and 90 mg of sterile lyophilized disodium pamidronate, each optionally containing 470 mg, 400 mg, and 375 mg of mannitol can readily be prepared and utilized as disclosed in Physician's Desk Reference (52d ed., 1988) at pages 1824-1828.

The present invention further provides an improved method for the preparation of aqueous solutions and dosage forms of disodium pamidronate of defined composition, using pamidronic acid as a starting compound and titrating an aqueous pamidronic acid slurry having a concentration of about 0.1 to 0.5 M, 10 preferably about 0.130 to about 0.135 M, to visual clarity at a pH of approximately 6.5 using aqueous sodium hydroxide (NaOH) or other suitable base to yield an aqueous solution of disodium pamidronate. Preferably, an about 2:1 molar ratio of NaOH or other metal hydroxide to pamidronic acid is used. Mannitol is an optional component of the slurry, preferably present in a molar excess to pamidronate as in the solutions described above, prior to lyophilization. Alternately, the solution is titrated to clarity using aqueous sodium hydroxide and the final pH adjusted to approximately  $6.5 \pm 0.1$  using phosphoric acid as needed. The clear aqueous solution of disodium pamidronate is then filtered and packaged as unit dosage forms comprising aliquots of the filtered solution.

This aspect of the present invention entirely eliminates the generation and isolation of various metastable, hydrated, crystalline and/or amorphous physical forms of disodium pamidronate (each of which can interchange with the other, depending upon the circumstance of available moisture or temperature in the substance environment). The present invention affords stable pamidronate compositions which can be stored at temperatures of, for example, 0°C or less (frozen), about 2 to about 8°C (refrigerated), or about 20 to about 30°C (room temperature). The compositions can be used to prepare dosage forms suitable for 30 therapeutic oral, rectal, transdermal, intranasal, or intravenous administration.

As used herein, the term "about" incorporates inherent variability of the parameter referenced, due to measurement techniques known to the art, i.e., pH, or water content (loss on drying).

20

25


4

## Detailed Description of the Invention

The present invention provides an improved method for the preparation of stable unit dosage forms of disodium pamidronate having constant composition. Pamidronic acid is used as a starting compound, from which an aqueous pamidronic acid slurry (the slurry being at least partially opaque) is formed. The concentration of pamidronic acid in the slurry is about 0.1 to 0.5 M, preferably about 0.130 to about 0.135 M. The pamidronic acid slurry is titrated to visual clarity, at a pH of approximately  $6.5 \pm 0.1$ , using aqueous sodium hydroxide (NaOH) or other suitable base to yield a solution of disodium pamidronate. Mannitol is an optional component of the slurry, and is preferably present in molar excess to the pamidronic acid. Alternately, the solution can be titrated to clarity using aqueous NaOH and the final pH can be adjusted to approximately 6.25 - 6.75 using phosphoric acid as needed.

The clear aqueous solution formed can be filtered and packaged as one or more unit dosage forms comprising aliquots of the filtered solution. The filtered solution can also be further processed by freezing, for example, at approximately -25 to -40°C and lyophilizing at 10-25 mbar at 20-40°C in situ to yield amorphous, essentially anhydrous (≤1-2 wt% water) disodium pamidronate which can be stored under nitrogen and readily reconstituted with sterile water or physiological salt solution for injection or infusion into a patient in need thereof. The invention described herein eliminates the need for the formation of metastable hydrates, which results in a dosage form of more constant composition than those described by Jary et al. (U.S. Patent No. 4,304,734) and Stahl et al. (U.S. Patent Nos. 4,711,880 and 4,639,338).

The solution of the present invention is used to prepare separately packaged oral or intravenous unit dosage forms. Unit dosage forms can comprise pamidronate solution prepared by the method of the present invention and packaged into pharmaceutical vials or ampules. Preferably, the containers are "single use" in that they contain one unit dosage form and are not reclosable. Also, it is preferably the employ container components that free of calcium (Ca<sup>+2</sup>) that can be sequestered by the pamidronate solution. Alternately, pamidronate solutions prepared by the method of the present invention can be used to prepare dosage forms such as, for example, soft gelatin capsules,



15

20

30


transdermal patches, intranasal sprays, rectal suppositories, or enteric-coated capsules.

Unit dosage forms can also comprise disodium pamidronate solution prepared by the method of the present invention wherein pamidronate is bound to one or more ion exchange resins. These compositions can be administered as, for example, liquid formulations or as sustained release dosage forms using polymer coatings known to those of skill in the art.

For intravenous administration, disodium pamidronate prepared by the method of the present invention can be packaged into vials or ampules comprising from 5 to 20 ml of the pamidronate solution, wherein the disodium pamidronate is present at a concentration of, for example, approximately 3 mg/ml to about 9 mg/ml. Prior to administration, disodium pamidronate solution within a vial or ampule is removed and diluted using an appropriate volume of a calcium-free solution such as a 0.9% w/v sodium chloride (normal saline) or a 5% dextrose solution.

For oral administration, capsule dosage forms can be prepared from anhydrous disodium pamidronate or from a disodium pamidronate solution prepared by the method of the present invention. Methods for preparing pharmaceutical formulations in capsule form are known to those of skill in the art. For example, one such method has been described in U.S. Patent No. 5,569,466 (Tanner et al.), which describes the preparation of fill compositions for soft gelatin capsules. Enteric coating compositions are also known to those of skill in the art. Enteric coatings are described, for example, in U.S. Patent No. 5,888,550 (Cook et al.), which describes cellulose acetate phthalate enteric coating compositions. With proper selection of polymer coatings, such as an acrylic-based resin designed to dissolve at a pH above 7, delivery to the distal ileum and colon can be achieved. An example of such an acrylic-based resin is the Eudragit E-3 (acrylate ester copolymer) series (Rohm-Haas, Philadelphia, PA).

U.S. Patent No. 5,780,055 (Habib et al.) describes beads of microcrystalline cellulose and a disintegrant for delivery of easy-to-swallow pharmaceutical compositions. U.S. Patent No. 5,431,920 (Bechard et al.) describes a double-coated bimodal release form of bisphosphonic acid for



25

administration to subjects exhibiting upper gastrointestinal tract sensitivity to bisphosphonic acid compounds. See, also, Strein (U.S. Patent No. 5,366,965). These methods, for example, can be used to prepare dosage forms of the disodium parnidronate solution of the present invention for administration to patients who, for example, have difficulty swallowing capsules or who exhibit gastric sensitivity to disodium parnidronate.

Methods for preparing suppository formulations are known to those of skill in the art and can be used to prepare suppository dosage forms comprising a disodium paraidronate or a disodium paraidronate solution in admixture with ingredients such as those described in U.S. Patent No. 4,439,194 (Harwood, et al.), which describes a water and drug delivery system for suppository use.

For intradermal delivery, transdermal patches can be formed using disodium pamidronate or a disodium pamidronate solution prepared by the method of the present invention. Methods for producing transdermal patches to deliver therapeutic compositions are described, for example, in U.S. Patent No. 5,932,240 (D'Angelo et al.), U.S. Patent No. 4,336,243 (Sanvordeker et al.), and U.S. Patent No. 5,650,165 (Akemi et al.). A transdermal patch or percutaneous absorption preparation can be held in contact with the skin by a pressure-sensitive adhesive. Examples of suitable pressure-sensitive adhesives include silicone pressure-sensitive adhesives and polyisobutylene pressure-sensitive adhesives, which are known to those of skill in the art. A pressure-sensitive adhesive suitable for use in forming a transdermal patch for disodium pamidronate delivery is described, for example, in U.S. Patent No. 5,785,985 (Czech et al.).

Formulations for intranasal delivery comprise, for example, nasal drops or sprays having a pamidronate concentration of approximately 3 mg/ml to about 30 mg/ml of carrier matrix. The invention will be further described by reference to the following detailed example.

The invention will be further described by reference to the following 30 detailed example.

15

7

#### Example 1

#### Preparation of Disodium Pamidronate

For a batch size of 5 L, 587.5 g (3.2 moles) of mannitol is dissolved in 3.5 L of water. Pamidronic acid (31.6 g, 0.133 moles) is mixed with a 1.0 L aliquot of the mannitol solution to form a slurry. The slurry is then transferred into the remainder of the mannitol solution, and stirred for at least 15 min. Aqueous 1 N sodium hydroxide (270 ml) is then added and the mixture is stirred until a clear, colorless solution results. The pH is then adjusted to 6.5 ± 0.1 using either 1 M aqueous phosphoric acid or 1 N aqueous sodium hydroxide, as needed. The solution is then filtered through a 0.22 micron filter, and filled at 20°C into vials at 4.0 ml (4.172 g)/vial, under sterile conditions. The aqueous solution is frozen at -37°C and lyophilized (20 mbar, 20-40°C) to yield 1,250 vials, each containing 30 mg of amorphous disodium pamidronate. The vials are sealed under positive nitrogen pressure. The disodium pamidronate is amorphous (noncrystalline) by X-ray diffraction and contains 0.7 wt-% water (USP 23 <731>).

#### Example 2

### Preparation of Disodium Pamidronate

For a batch size of 5 L, 587.5 g (3.2 moles) of mannitol is dissolved in

3.5 L of water. Pamidronic acid (31.6 g, 0.133 moles) is mixed with a 1.0 L
aliquot of the mannitol solution to form a slurry. The slurry is transferred into
the remainder of the mannitol solution, and stirred for about 15 min. Aqueous
1 N sodium hydroxide (270 ml) is added and the mixture is stirred until a clear,
colorless solution results. The pH is adjusted to 6.5 ± 0.1 using either 1 M


25 aqueous phosphoric acid or 1 N aqueous sodium hydroxide, as needed. The
solution is filtered through a 0.22 micron filter, and filled at 20°C into glass vials
at 4.0 ml (4.172 g)/vial, and the vials are sealed in the lyophilizer in an inert
atmosphere under sterile conditions.

All publications, patents and patent documents are incorporated by reference herein, as though individually incorporated by reference. The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many

WO 00/34293


8

variations and modifications may be made while remaining within the spirit and scope of the invention.



#### WHAT IS CLAIMED IS:

- 1. A method for preparing disodium pamidronate comprising:
  - (a) preparing a stirred slurry of pamidronic acid in water;
- 5 (b) adding aqueous solution of sodium hydroxide to said slurry in an about 2:1 molar ratio of sodium hydroxide to pamidronic acid; to yield a clear solution having a pH of about 6.5;
  - (c) filtering said solution; and
- (d) freezing and lyophilizing said filtered solution to yield amorphous, essentially annydrous disodium pamidronate.
  - 2. The method of claim 1 wherein said slurry comprises mannitol.
- 3. The method of claim 1 or 2 wherein, in step (d) a plurality of aliquots of solution are filled into a plurality of storage vessels, frozen and lyophilized in situ.
- The method of claim 3 wherein the amorphous, essentially anhydrous disodium pamidronate is sealed into said vessels under positive dry nitrogen
   pressure.
  - 5. The method of claim 1 wherein the essentially anhydrous disodium pamidronate contains ≤ 2 wt-% water.
- 25 6. The method of claim 1 wherein the solution is frozen at about -25°C to -40°C
  - 7. The method of claim 1 wherein the lyophilization is carried out at about 20-40°C at about 10-25 mbar.
  - 8. Disodium pamidronate prepared by the process of claim 1.



10

20

25

- 9. A method for preparing a therapeutic aqueous disodium pamidronate solution comprising:
  - (a) preparing a slurry of pamidronic acid in water, and
- (b) combining aqueous sodium hydroxide with said slurry in an about 2:1 molar ratio of sodium hydroxide to pamidronic acid; to yield a solution of disodium pamidronate having visual clarity and a pH of about 6.5;
  - (c) packaging said solution in a plurality of sealed containers to yield a plurality of liquid unit dosage forms of pamidronate.
- 10. The method of claim 9 wherein the slurry includes an effective stabilizing amount of mannitol.
- 11. A unit dosage form comprising a disodium pamidronate solution prepared by the method of claim 9 or 10.
  - 12. A capsule comprising a unit dosage of the solution of claim 11.
  - 13. The capsule of claim 12 which comprises an enteric coating.
  - 14. An intranasal spray dispenser comprising at least one unit dosage form of the solution of claim 11.
  - 15. A vial or ampule comprising a unit dosage form of the solution of claim
    11.
    - 16. The vial or ampule of claim 15 wherein the solution is packaged under an inert atmosphere.
- 30 17. The dosage form of claim 11 wherein the container is free of Ca<sup>+2</sup> that can be sequestered by the disodium pamidronate.

- 18. The dosage form of claim 17 wherein the container is a plastic vial or ampule.
- 19. The dosage form of claim 17 wherein the container is a glass vial or5 ampule.

## INTERN OI SEARCH REPORT

nel Application No.

PCT/US 99/29284

A CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07F9/38 A61K31/663

A61K47/26

A61K9/4B

A61K9/12

According to International Patent Classification (IPC) or to both national classification and IPC

#### B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symposs) IPC 7 CO7F A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category *	Chatton of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A	EP 0 177 443 A (CIBA-GEIGY AG) 9 April 1986 (1986-04-09) cited in the application the whole document	1–19		
<b>A</b>	CHEMICAL ABSTRACTS, vol. 128, no. 10, 9 March 1998 (1998-03-09) Columbus, Ohio, US; abstract no. 115081, SU G ET AL: "Preparation of amino-diphosphinic acid and its sodium salts as bone absorption inhibitors" XP002134859 abstract & CN 1 144 806 A (NANJING PHARMACEUTICALS INST.; PEOP. REP. CHINA)	1-19		

Further documents are listed in the continuation of box C.	Patent tamily members are listed in annex.		
*Special categories of cited documents :  *A* document defining the general state of the last which is not considered to be of particular relevance.	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention		
*E" earlier document but published on or after the international filing date  "L" document which may throw doubts on priority claim(a) or	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone. "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.		
which is cited to establish the publication date of another citation or other special reason (as apecified)  "O" document reterring to an oral disclosure, use, exhibition or other meens  "P" document published prior to the international filing date but			
later than the priority date claimed  Date of the actual completion of the international search	"&" document member of the same patent family  Date of making of the international search report		
10 April 2000	25/04/2000		
Name and mailing address of the ISA	Authorized officer		
European Patent Office, P.B. 5616 Patentiaan 2 NL – 2280 HV Rijswijk Tel. (≼31–70) 340–2040, Tx. 31 651 epo nl, Fax: (≼31–70) 340–3016	Beslier, L		

1



# SEARCH REPORT

4			
	Licen and	Application No.	
	PCT/US	99/29284	

C.(Continu	C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication,where appropriate, of the resevant passages	Relevant to claim No.				
P,A	DE 198 20 974 A (HENKEL KGAA ) 18 November 1999 (1999-11-18) the whole document	1-19				

1

# INTERNATION EARCH REPORT

PCT/US 99/29284

enformation on patent tamby members

Patent document cited in search report		Publication date :	Patent tamily member(s)	Publication date
EP 177443	A	09-04-1986	AR 240832 A	28-02-1991
El 1//440	••		AT 47396 T	15-11-1989
			AU 591066 B	30-11-1989
			AU 4577785 A	13-02-1986
			CA 1255694 A	13-06-1989
			CY 1570 A	20-12-1991
			DD 239597 A	01-10-1986
			DK 355485 A	07-02-1986
			ES 545899 D	16-11-1986
			ES 8701188 A	16-02-1987
			ES 555602 D	01-07-1987
			ES 8706707 A	16-09-1987
			FI 852986 A,B	, 07-02-1986
			GR 851910 A	03-12-1985
			HK 42591 A	07-06-1991
			HU 38361 A,E	28-05-1986
			IE 58077 B	30-06-1993
			IL 75990 A	31-07-1989
J			JP 1797264 C	28-10-1993
			JP 5008717 B	02-02-1993
			JP 61043196 A	01-03-1986
			KR 9400817 B	02-02-1994
			MX 9203373 A	01-09-1992
			NO 853087 A.E	3, 07-02-1986
			NO 902523 A,	3, 07-02-1986
			PH 24854 A	26-12-1990
			PT 80916 A,I	3 01-09-1985
			SG 91390 G	18-01-1991
			US 4639338 A	27-01-1987
			US 4711880 A	08-12-1987
			ZA 8505892 A	26-03-1986
CN 1144806	A	12-03-1997	NONE	
DE 19820974	A	18-11-1999	NONE	